

## Key to the subtribes

- 1 Inner eye margins distinctly converging above . . . . . Heliocausina
- Inner eye margins nearly parallel or converging toward clypeus . . . . . 2
- 2 Posterolateral oblique scutal carina absent . . . . . Exeirina
- Posterolateral oblique scutal carina present . . . . . 3
- 3 Scutellum not overlapping metanotum . . . . . Gorytina
- Scutellum overlapping metanotum . . . . . 4
- 4 Forewing stigma moderately large . . . . . Spheciina
- Forewing stigma minute . . . . . 5
- 5 Forewing submarginal cell I with prestigmal length not more than length of wing beyond marginal cell. . . . . Handlirschiina
- Forewing submarginal cell I with prestigmal length much more than length of wing beyond marginal cell . . . . . 6
- 6 Labrum prominent, exposed part nearly half as long as broad . . . . . Stizina
- Labrum markedly prominent, exposed part at least as long as broad . . . . . 6
- 7 Ocelli depressed below level of surrounding surface of vertex; midocellar lens present, elongate, oval, sometimes medially indented . . . . . Stictiellina
- Ocelli not depressed below level of surrounding surface of vertex; midocellar lens reduced to a scar, at most with traces of a lens . . . . . Bembicina

Extract from:

Nemkov, P.G. & Ohl, M. 2011. A cladistic analysis and reclassification of the tribe Bembicini (Hymenoptera: Crabronidae: Bembicinae). *Zootaxa*, 2801: 27-47

